

AN OPTIMAL LOWER CURVATURE BOUND FOR CONVEX HYPERSURFACES IN RIEMANNIAN MANIFOLDS

STEPHANIE ALEXANDER, VITALI KAPOVITCH, AND ANTON PETRUNIN

ABSTRACT. It is proved that a convex hypersurface in a Riemannian manifold of sectional curvature $\geq \kappa$ is an Alexandrov's space of curvature $\geq \kappa$. This theorem provides an optimal lower curvature bound for an older theorem of Buyalo.

The purpose of this paper is to provide a reference for the following theorem:

Theorem 1. *Let M be a Riemannian manifold with sectional curvature $\geq \kappa$. Then any convex hypersurface $F \subset M$ equipped with the induced intrinsic metric is an Alexandrov's space with curvature $\geq \kappa$.*

Here is a slightly weaker statement:

Theorem 2. [Buyalo] *If M is a Riemannian manifold, then any convex hypersurface $F \subset M$ equipped with the induced intrinsic metric is locally an Alexandrov's space.*

In the proof of Theorem 2 in [Buyalo], the (local) lower curvature bound depends on (local) upper as well as lower curvature bounds of M . We show that the approach in [Buyalo] can be modified to give Theorem 1.

Definition 3. *A locally Lipschitz function f on an open subset of a Riemannian manifold is called λ -concave ($\lambda \in \mathbb{R}$) if for any unit-speed geodesic γ , the function*

$$f \circ \gamma(t) - \lambda t^2 / 2$$

is concave.

Lemma 4. *Let $f : \Omega \rightarrow \mathbb{R}$ be a λ -concave function on an open subset Ω of a Riemannian manifold. Then there is a sequence of nested open domains Ω_i , with $\Omega_i \subset \Omega_j$ for $i < j$ and $\cup_i \Omega_i = \Omega$, and a sequence of smooth λ_i -concave functions $f_i : \Omega_i \rightarrow \mathbb{R}$ such that*

- (i) *on any compact subset $K \subset \Omega$, f_i converges uniformly to f ;*
- (ii) *$\lambda_i \rightarrow \lambda$ as $i \rightarrow \infty$.*

This lemma is a slight generalization of [Greene–Wu, Theorem 2] and can be proved exactly the same way.

Proof of Theorem 1. Without loss of generality one can assume that

- (a) $\kappa \geq -1$,

1991 *Mathematics Subject Classification.* Primary 53C20, 53B25; Secondary 53C23, 53C45.

We thank the Mathematical Research Institute Oberwolfach (Research in Pairs programme) for a valuable visit, of which this note is an offshoot. The third-named author was supported in part by the National Science Foundation under grant DMS-0406482.

- (b) F bounds a compact convex set C in M ,
- (c) there is a (-2) -concave function μ defined in a neighborhood of C and $|\mu(x)| < 1/10$ for any $x \in C$,
- (d) there is unique minimal geodesic between any two points in C .

(If not, rescale and pass to the boundary of the convex piece cut by F from a small convex ball centered at $x \in F$, taking $\mu = -10 \operatorname{dist}_x^2$.)

Consider the function $f = \operatorname{dist}_F$. By Rauch comparison (as in [Petersen, 11.4.8]), for any unit-speed geodesic γ in the interior of C , $(f \circ \gamma)''$ is bounded in the support sense by the corresponding value in the model case (where $M = \mathbb{H}^2$ and F is a geodesic). In particular,

$$(f \circ \gamma)'' \leq f \circ \gamma.$$

Therefore $f + \varepsilon \mu$ is $(-\varepsilon)$ -concave in $\Omega_\varepsilon = C \cap f^{-1}((0, \varepsilon))$. Take $K_\varepsilon = f^{-1}([\frac{1}{3}\varepsilon, \frac{2}{3}\varepsilon]) \cap C$. Applying lemma 4, we can find a smooth $(-\frac{\varepsilon}{2})$ -concave function f_ε which is arbitrarily close to $f + \varepsilon \mu$ on K_ε and which is defined on a neighborhood of K_ε . Take a regular value $\vartheta_\varepsilon \approx \frac{1}{2}\varepsilon$ of f_ε . (In fact one can take $\vartheta_\varepsilon = \frac{1}{2}\varepsilon$, but it requires a little work.) Since $|\mu|_C < 1/10$, the level set $F_\varepsilon = f_\varepsilon^{-1}(\vartheta_\varepsilon)$ will lie entirely in K_ε . Therefore F_ε forms a smooth closed convex hypersurface. By the Gauss formula, the sectional curvature of the induced intrinsic metric of F_ε is $\geq \kappa$. F_ε bounds a compact convex set C_ε , where $F_\varepsilon \rightarrow F$, $C_\varepsilon \rightarrow C$ in Hausdorff sense as $\varepsilon \rightarrow 0$. By property (d), the restricted metrics from M to C, C_ε are intrinsic, and so C_ε is an Alexandrov space with F_ε as boundary, that converges in Gromov–Hausdorff sense to C . It follows from [Petrinin, Theorem 1.2] (compare [Buyalo, Theorem 1]) that F_ε equipped with its intrinsic metric converges in Gromov–Hausdorff sense to F equipped with its intrinsic metric. Therefore F is an Alexandrov space with curvature $\geq \kappa$. \square

Remark 5. We are not aware of any proof of theorem 1 which is not based on the Gauss formula. (Although if M is Euclidean space, there is a beautiful purely synthetic proof in [Milka].) Finding such a proof would be interesting on its own, and also could lead to the generalization of theorem 1 to the case when M is an Alexandrov space.

REFERENCES

- [Buyalo] Buyalo, S., Shortest paths on convex hypersurface of a Riemannian manifold (Russian), Studies in Topology, Zap. Nauchn. Sem. LOMI 66 (1976) 114–132; translated in J. of Soviet Math. 12 (1979), 73–85.
- [Greene–Wu] Greene, R. E.; Wu, H.-H., On the subharmonicity and plurisubharmonicity of geodesically convex functions. Indiana Univ. Math. J. 22 (1972/73), 641–653.
- [Milka] Milka, A. D. Shortest arcs on convex surfaces (Russian), Dokl. Akad. Nauk SSSR 248 1979, no. 1, 34–36; translated in Soviet Math. Dokl. 20 (1979), 949–952.
- [Petersen] Petersen, P., Riemannian Geometry, Springer-Verlag, New York Berlin Heidelberg, 1998.
- [Petrinin] Petrunin, A., Applications of quasigeodesics and gradient curves. Comparison geometry (Berkeley, CA, 1993–94), 203–219, Math. Sci. Res. Inst. Publ., 30, Cambridge Univ. Press, Cambridge, 1997.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, IL 61801

E-mail address: sba@math.uiuc.edu

URL: www.math.uiuc.edu/~sba

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ON, M5S 2E4

E-mail address: vtk@math.toronto.edu

URL: www.math.toronto.edu/~vtk/

DEPARTMENT OF MATHEMATICS, PENN STATE UNIVERSITY, STATE COLLEGE, PA 16802

E-mail address: petrunin@math.psu.edu

URL: www.math.psu.edu/petrunin/